

## POLYMERS

**Publication number:** JP2001508480T

**Publication date:** 2001-06-26

**Inventor:**

**Applicant:**

**Classification:**




- international: **A61L27/00; A61F9/007; C08F220/18; C08F220/28; C08F220/58; C08F230/02; C08F230/08; C09D143/04; A61L27/00; A61F9/007; C08F220/00; C08F230/00; C09D143/00;** (IPC1-7): C08F220/18; A61L27/00; C08F220/28; C08F230/02; C09D143/04

- european: A61F9/007V; C08F220/28; C08F220/58; C08F230/08

**Application number:** JP19980530659T 19980112

**Priority number(s):** GB19970000390 19970110; GB19970018909 19970905; WO1998GB00070 19980112

**Also published as:**

 WO9830615 (A1)  
 EP0953001 (A1)  
 EP0953001 (A0)

**Report a data error here**

Abstract not available for JP2001508480T

Abstract of corresponding document: **WO9830615**

A cross-linkable polymer is formed from monomers including a) a hydroxyl-group containing monomer of general formula (I) in which R<26> is hydrogen or C1-4-alkyl group, A<1> is -O- or -NR<27>- where R<27> is hydrogen or a C1-4-alkyl group or a group B<9> R-(OH)<sub>n</sub>; B<9> is a bond, a straight or branched alkylene, an oxaalkylene or oligooxa alkylene group; R<13> is a n + 1 functional (optionally substituted) C1-24 alkylene group, n is an integer of one or more; and b) a reactive monomer of general formula (II) in which R<19> is hydrogen or a C1-4-alkyl group or A<2> is -O- or -NR<21>- where R<21> is hydrogen, or a C1-4-alkyl group, R<12> is C1-24 straight or branched alkylene, oxaalkylene or oligo oxaalkylene group in which each alkylene group has 1 to 6 carbon atoms, A<3> is a band or -O-, each R<16> independently selected from C1-6-alkyl groups. Preferably the polymer includes zwitterionic monomer and is useful for providing stable biocompatible coatings on substrates.

---

Data supplied from the **esp@cenet** database - Worldwide